

CS 3 SEM APPLIED MATHEMATICS-III JUN 2016

QP Code : 30557

(3 Hours)

[Total Marks : 80]

- 1) Question No. 1 is compulsory.
- 2) Attempt any THREE of the remaining.
- 3) Figures to the right indicate full marks.

-
- Q 1.A) If $\int_0^{\infty} e^{-2t} \sin(t + \alpha) \cos(t - \alpha) dt = \frac{1}{4}$, find α (5)
- B) Find half range Fourier cosine series for $f(x) = x$, $0 < x < 3$. (5)
- C) If $u(x,y)$ is a harmonic function then prove that $f(z) = u_x - iu_y$ is an analytic function. (5)
- D) Prove that $\nabla f(r) = f'(r) \frac{r}{r}$ (5)
- Q.2) A) If $v = e^x \sin y$, prove that v is a harmonic function. Also find the corresponding analytic function. (6)
- B) Find Z-transform of $f(k) = b^k$, $k \geq 0$ (6)
- C) Obtain Fourier series for $f(x) = \frac{3x^2 - 6x\pi + 2\pi^2}{12}$ in $(0, 2\pi)$,
where $f(x+2\pi) = f(x)$. Hence deduce that $\frac{\pi^2}{6} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots$ (8)
- Q.3) A) Find inverse Laplace of $\frac{(s+3)^2}{(s^2+6s+5)^2}$ using Convolution theorem (6)
- B) Show that the set of functions $\{\sin x, \sin 3x, \sin 5x, \dots\}$ is orthogonal over $[0, \pi/2]$. Hence construct orthonormal set of functions (6)
- C) Verify Green's theorem for $\int_C \frac{1}{y} dx + \frac{1}{x} dy$ where C is the boundary of region defined by $x = 1, x = 4, y = 1$ and $y = \sqrt{x}$ (8)

TURN OVER

- 2 -

Q.4) Find $Z\{k^2 a^{k-1} U(k-1)\}$ (6)

B) Show that the map of the real axis of the z-plane is a circle under the transformation $w = \frac{2}{z+i}$. Find its centre and the radius. (6)

C) Express the function $f(x) = \begin{cases} \sin x & |x| < \pi \\ 0 & |x| > \pi \end{cases}$ as Fourier sine Integral. (8)

Q.5) A) Using Gauss Divergence theorem evaluate $\iint_S \bar{N} \cdot \bar{F} ds$

where $\bar{F} = x^2 i + zj + yzk$ and S is the cube bounded by $x = 0, x = 1, y = 0, y = 1, z = 0, z = 1$ a2zSubjects.com (6)

B) Find inverse Z-transform of $F(z) = \frac{z}{(z-1)(z-2)}$, $|z| > 2$ a2zSubjects.com (6)

C) Solve $(D^2 + 3D + 2)y = e^{-2t} \sin t$, with $y(0) = 0$ and $y'(0) = 0$ (8)

Q.6) A) Find Fourier expansion of $f(x) = 4 - x^2$ in the interval $(0,2)$ (6)

B) A vector field is given by $\bar{F} = (x^2 + xy^2) i + (y^2 + x^2y) j$. Show that \bar{F} is irrotational and find its scalar potential. (6)

C) Find (i) $L^{-1}\left\{\tan^{-1}\left(\frac{x}{s}\right)\right\}$

(ii) $L^{-1}\left(\frac{e^{-ms}}{s^2 - 2s + 2}\right)$ (8)

---XXX---