5

5

6

6

- N.B.: (1) Questions No. 1 is compulsory.
 - (2) Attempt any three questions from the remainirig five questions.
 - (3) Figures to the right indicate full marks.
- (a) Find half range sine series for $f(x) = x \sin x$ in $(0, \pi)$.
 - (b) Find the work dene is moving a pacticle once round the ellipse the plane z = 0 in the four field given by $\overline{F} = (3x - 2y)i + (2x + 3y)j + y^2k$.
 - (c) Evaluate $\int \overline{F} \cdot dr$ where $\overline{F} = yzi + zxj + xy$ kand C is the portion of the curve. 5
 - $\bar{r} = a \cos t i + b \sin t j + ct k$ from t = 0 to $t = \pi/4$
 - (d) Show that the set of funtion $f(r) = \sin((2n+1)x)$; n = 0, 1, 2, 3--- in $(0, \pi/2)$ is orthogonal. Hence construct the orthonormal set of functions.
- (a) Find the Fourier series of $f(x) = \frac{\pi x}{\pi}(2 x)$; $0 \le x \le 1$
 - (b) Evaluate ∫∫(∇xF)-ds where

 $\vec{F} = (2x - y + z) i + (x + y - z^2) + (3x - 2y + 4z) k$ and S is the surface of the cylinder $x^2 + y^2 = 4$ bounded by the plane z = 9 and open at the other end.

(c) A tightly stretched string with fixed end points x = 0 and x = l, in the shape defined by y = kx ($(x - \bar{x})$) where k is a constant, is released from this position of rest. Find y(x, t), the vertical displacement if

$$\frac{\partial^2 y}{\partial t^2} = C^2 \frac{\partial^2 y}{\partial x^2}.$$

(a). An eleastic string is stretched between two points at a distance π apart. In its equillibrium position the string in the shape of the curve $f(x) = K(\sin x - \sin 2x)$. Obtain y(x, t) the vertical displacement if y satisfies the equation

$$\frac{\partial^2 y}{\partial t^2} = \frac{\partial^2 y}{\partial x^2}.$$

[TURN OVER

GN-Con.:8584-14.

(b) Find Fourier integral representation of

$$f(x) = \frac{e^{ax}}{e^{-ax}}; x \le 0$$
; $a > 0$.

Hence show that

$$\int_{0}^{\infty} \frac{\cos \omega x}{\omega^{2} + a^{2}} d\omega = \frac{\pi}{2a} e^{-ax}; x > 0, a > 0.$$

(c) Solve the equation $\frac{\partial u}{\partial t} = K \frac{\partial^2 u}{\partial x^2}$ for the conduction of heat along a rod without

radiation subject to the following conditions:-

- (i) u is finite when t→∞ muADDA.com
- (ii) $\partial u/\partial x = 0$ when x = 0 for all values of t.
- (iii) u = 0 when $x = \ell$ for all values of t.
- (iv) $u = u_0$ when t = 0 for $0 < x < \ell$.
- 4. (a) Evaluate $\iint \overline{F} d\overline{s}$ where $\overline{F} = 4xi 2y^2 j + 2^2 K$ and S is the region bounded by 6

$$y^2 = 4x$$
, $x = 1$, $z = 0$, $z = 3$.

- $y^2 = 4x$, x = 1, z = 0, z = 3. (b) Find Fourier series for $f(x) = |\sin x|$ in $(-\pi, \pi)$.

$$f(x) = \frac{3x^2 - 6x\pi + 2\pi^2}{12}$$
 in $(0, 2\pi)$. Hence declare that

$$\frac{\pi^2}{6} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

- Obtain the complex form of Fourier series for $f(x) = \cosh x$ is $(-\ell, \ell)$.
 - (b) Obtain a solution of $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ to satisfy the following conditions
 - (i) $u \to 0$ as $y \to \infty$ for all x.
 - (ii) u = 0, if x = 0 for all y. muADDA.com
 - (iii) u = 0 if $x = \ell$ for all y
 - (iv) $u = \ell x x^2$ if y = 0 for all values of x between 0 and ℓ .
 - (c) If the vector field F is irrotational find the constants a, b, c where F is given by $\overline{F} = (x + 2y + az) i + (bx - 3y - z) j + (4x + cy + 2z) K$ Find the scalar potential for F and also find the work done in moving a practicle in this field from (1, 2, -4) to (3, 3, 2) along the straight line joining these points.

[TURN OVE

6 8

nuADDA.com

GN-Con.:8584-14.

3

QP Code: 12455

(a) Find Fourier sine integral representation for

6

$$f(x) = \frac{e^{-ax}}{x} : x > 0.$$

(b) Evaluate by Green's Theorem

$$\int_{c} (3x^2 - 8y^2) dx + (4y - 6xy) dy$$

where C is the boundary of the region bounded by $y = x^2$ and $y^2 = x$.

(c) Optain Fourier series for

8

$$f(r) = x + \frac{\pi}{2} - \pi < x < 0$$

$$\frac{\pi}{2} - x \quad ; 0 < x < \pi$$

Hence deduce that

$$\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$$

GN-Con.:8584-14