

(1) 26
(REVISED COURSE)
(3 Hours)

Comp Programming & numerical
CHEM/CBGS/III/CINM
QP Code :4908

[Total Marks: 80]

N.B. :

- 1) Question - 1 is compulsory. Answer any three questions from remaining.
- 2) Assume data if necessary and specify the assumptions clearly
- 3) Draw neat sketches wherever required.
- 4) Answer to the sub-questions of an individual question should be grouped and written together i.e. one below the other.

1. (a) Consider the following system of equations: [10]

$$\begin{bmatrix} 1 & 1 \\ 1 & (1+\epsilon) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

Solve the system for $\epsilon = 0.01$; 0.001 ; and 0.0001 by elimination. How sensitive is the solution to changes in ϵ ?

- (b) Use Crank-Nicholson Scheme to solve, [05]

$$u_{xx} = u_t \quad 0 \leq x \leq 1 \quad t > 0 \quad h = \frac{1}{4} \quad k = \frac{1}{4}$$

Given $u(x, 0) = 0$, $u(0, t) = 0$, $u(1, t) = 50t$ Compute u for one step in t -direction.

- (c) Explain how to use 'for' loop in SciLab with appropriate example. [05]

2. (a) Consider the following equation: [12]

$$f(x) = x^3 - 5x^2 + 6x - 1 = 0$$

Rewriting this as:

$$x = x + f(x) = g(x)$$

Start from $x^{(0)} = 0$ and compute upto $x^{(3)}$, by Successive Substitution. What happens if we start with $x^{(0)} = 1$. Is convergence guaranteed, for $0 \leq x \leq 3$?

- (b) Solve the following system by LU decomposition: [08]

$$\begin{bmatrix} 1 & -1 & 2 \\ 2 & -2 & 3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -6 \\ -20 \\ -2 \end{bmatrix}$$

3. (a) Obtain a root of the following equation using Newtons Method in the interval [12]
 $0 \leq x \leq 2$:

$$f(x) = \ln(x^2 + 1) - e^{0.4x} \cos(\pi x)$$

- (1) Solve the following equation to find a root in the interval $[0.1, 2]$ by the method [08]
false position. Obtain a value correct to three decimal places.

4. (a) Find $x(2)$ from following ODE using fourth order Runge-Kutta method: [14]

$$\frac{dx}{dt} = e^{-t} (\cos t - \sin t) - \sin\left(\frac{t}{2}\right) \quad \text{with} \quad x(0)$$

26/5/15

(2)

Chem/CBGS/III

2

QP Code :4908

- (b) Write Laplace equation and express it in difference form using Taylor's series expansion [06]

5. Solve the following equations using Newtons Method and starting from [20]

$$x^{(0)} = \begin{bmatrix} 0.1 \\ 0.2 \end{bmatrix}$$

$$f_1(x) = x_1^2 x_2 + x_2^2 x_1 - 2x_1 x_2 + 5x_1 - 7 = 0$$

$$f_2(x) = x_1^3 - 2x_1^2 x_2 + x_1 x_2^2 - 6x_2 + 15 = 0$$

Perform minimum three iterations.

5. (a) Consider the following example: [12]

$$\frac{dy}{dt} = -y^2 \quad \text{with} \quad y = 1 \quad \text{at} \quad t = 0$$

where t and y are scaled variables. This equation represents a second order reaction taking place in a tubular reactor. Integrate the equation using Eulers Implicit Method from $t = 0$ to $t = 5$.

- (b) Solve the following system by Jacobi Iteration: [08]

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & -2 & 1 & 0 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

Carry out five iterations.

JF-Con.: 19633-15.