Chemical Engineering Thermodynamics-I

CHEM/17/CBGS/CET-1

(3 Hours)

[Total Marks: 100

N.B.: (1) Question No. 1 is compulsory.

- (2) Attempt any Three of the remaining questions.
- Each question carries Equal Marks.
- l. Solve any Four of the following.
 - 5 Three moles of nibgen at 30°C contained in a rigid vessel, its heated to 250°C. How much heat us required to do this, it vessel weights 100 kg and has a capacity of 0.5 $\frac{KJ}{kg^{\circ}C}$, how much heat is required? For nitrogen $C_v = 20.8 \text{ J/mol}^{\circ}C_v = 29.1 \text{ J/mol}^{\circ}C.$
 - (b) Give statement of first law of thermodynamics and its mathematical form 5 when applied to different processes.
 - A carnot engine operating between 800°C and 25°C is used to run a carnot refrigerator operating between -20°C and 25°C. It the engine aborhas 10 KJ/s from the reservoir at 800°C, determine the capacity of the refrigerator.
 - Define fugacity and fugacity coefficient.

- 5
- What is a difference between state function and path function.
- 5
- 2. One kmol of an ideal gas at 100 kPa and 300K undergoes the following reversible 20 changes.
 - (i) Compressed adiabatically to 500 kPa.
 - (ii) Heated at constant pressure to 800 K.
 - (iii) Expanded adiabatically to 210 kPa.
 - (iv) Cooled at constant volume to 300 K.
 - Expanded isothermally to 100 kPa.

Find AH, Q, AU & W for the individual stage and also for the entire cycle.

Also find the thermal efficiency of the process.

Cp = 29.099 kJ/kmol k, Cv = 20.785 kJ/kmol K.

- Find the volume of n-pentane at 500 K and 20 bar for the following cases: 10 3.
 - (i) As an ideal gas.
 - (ii) As Van der waals gas. Tc = 469.6 K, Pc = 33.7 bar
 - For an adiabatic process prove that (b)

10

$$W = \frac{P_1 V_1 - P_2 V_2}{v - 1}$$

10

27/5/15



CHEM/ W/CBGS/CET-1

Q.P. Code: 3633

2

4. (a) Find the compressibility foctor at the critical point for a gas obeying vender 10 waal equation of state.

$$\left(z + \frac{27}{64} \frac{Pr}{ZTr^2}\right) \left(1 - \frac{Pr}{8ZTr}\right) = 1$$

- (b) A vessel is divided into two parts by a partition, on one side 4 kmol of nitrogen gas at 80°C and 40 bar and on the other side 2 kmol of argon at 120°C and 20 bar are kept. If the partition is removed and the gases are mixed adiabatically, what is the change in entropy? Assume N₂ as an ideal gas. Cp = 5/2 R, Cv = 3/2R.
- 5. (a) Explain the concept of exergy and get the expression to calculate exergy 10 loss when system changes its state.
 - (b) Calculate the enthalpy and entropy departure for n-octance vapor at 427.85 K and 0.215 MPa, using the generalized Redlich-kwong equation of state a = 4.426 m⁶Pa Mol² and b = 164.3x10⁻⁶m³/mol; Z = 0.9308, B=9.9306x10⁻³.
- 6. (a) Derive Maxwells equations.
 - (b) Write note on:
 - (b) Write note on:(i) Clauses Inequality.
 - (i) Clauses Inequality.(ii) Joule Thompson Coefficient.

JP-Con. 10861-15.